Brain Research, 478 (1989) 407–409
Elsevier

Author Index

Abate, C., see Milner, T.A., 1
Ableitner, A. and Herz, A., Limbic brain structures are important sites of κ-opioid receptor-mediated actions in the rat: a [14C]-2-deoxyglucose study, 326
Adret, P. and Rogers, L.J., Sex difference in the visual projections of young chicks: a quantitative study of the thalamofugal pathway, 59
Alvarez, J., see Von Bernhardi, R., 301
Amir, S., Pollock, R. and Lagiorgia, M., Intra-hypothalamic injection of thyrotropin-releasing hormone suppresses brown fat thermogenesis in the anaesthetized rat, 361
Anderson, J., Wallace, W., Snyder, S., Haroutunian, V., Roberts, J.L. and Lieberburg, I., Cellular forms of the rat and human β-amyloid precursor protein (BAPP), 391
Andiné, P., see Kjellmer, I., 241
Ashida, H., see Maru, E., 112

Baumann, N., see Bourdette, D.N., 175
Bennett, B., see MacLean, D.B., 349
Benson, D.M., Blitzer, R.D., Haroutunian, V. and Landau, E.M., Functional muscarinic supersensitivity in denervated rat hippocampus, 399
Bermúdez-Rattoni, F., see Escobar, M., 368
Bérod, A., see Kitahama, K., 156
Blitzer, R.D., see Benson, D.M., 399
Boeijinga, P.H. and Lopes da Silva, F.H., Modulations of EEG activity in the entorhinal cortex and forebrain olfactory areas during odour sampling, 257
Boggan, W.O. and Ondo, J.G., Effect of central administration of phencyclidine on plasma concentrations of luteinizing hormone, 382
Bommelaer-Bayt, M.-C., see Watanabe, Y., 143
Bourdette, D.N., Zalc, B., Baumann, N. and Seil, F.J., Antisera to the ganglioside GM1 do not have anti-myelin or anti-axon activities in vitro, 175
Bourret, C. and Mallart, A., Depression of calcium current at mouse motor nerve endings by polycationic antibiotics, 403
Bowersox, J., see Starkstein, S.E., 16
Britton, D.R. and Indyk, E., Effects of ganglionic blocking agents on behavioral responses to centrally administered CRF, 205
Brot, M., see Shewey, L.M., 95
Butler, B., see Gunion, M.W., 47

Carpenter, C.L., see Marks, S.S., 196
Casper, D. and Davies, P., Mechanism of activation of choline acetyltransferase in a human neuroblastoma cell line, 85
Casper, D. and Davies, P., Stimulaton of choline acetyltransferase activity by retinoic acid and sodium butyrate in a cultured human neuroblastoma, 74
Chiang, C.-Y. and Zhou, M., Evidence for the involvement of a descending cholinergic pathway in systemic morphine analgesia, 293
Chin, G.J., see Vogel, S.S., 281
Chopp, M., see Mesina, J.E., 181
Clementi, G., Favit, A., Curreri, R., Nicoletti, F. and Scapagnini, U., Repeated calcitonin treatment reduces the stimulation of inositol phospholipid by norepinephrine and serotonin in rat hippocampus and cerebral cortex, 138
Colby, K.A., Thompson, T.L. and Patrick, R.L., Tyrosine hydroxylase phosphorylation in rat brain striatal synaptosomes, 103
Curreri, R., see Clementi, G., 138

Davies, P., see Casper, D., 74
Davies, P., see Casper, D., 85
De Bruyne, K., see Gerstner, G.E., 233
De Saint Hilaire-Kafi, Z., Gibbs, J. and Nicolaidis, S., Satiety and sleep: the effects of bombesin, 152
Dorsa, D.M., see Shewey, L.M., 95
Dray, F., see Watanabe, Y., 143
Dusart, I., see Juliano, S.L., 356

Eldridge, J.C., Fleenor, D.G., Kerr, D.S. and Landfield, P.W., Impaired up-regulation of type II corticosteroid receptors in hippocampus of aged rats, 248
Emerich, D.F., see Walsh, T.J., 24
Eriksdotter-Nilsson, M., Gerhardt, G., Seiger, Å., Olson, L., Hoffer, B. and Granholm, A.-C., Age-related alterations in noradrenergic input to the hippocampal formation: structural and functional studies in intraocular transplants, 269
Escobar, M., Fernández, J., Guevara-Aguilar, R. and Bermúdez-Rattoni, F., Fetal brain grafts induce recovery of learning deficits and connectivity in rats with gustatory neocortex lesion, 368

Faraci, F.M., Mayhan, W.G. and Heistad, D.D., Effect of serotonin on blood flow to the choroid plexus, 121
Favit, A., see Clementi, G., 138
Fernández, J., see Escobar, M., 368
Fischer-Colbrie, R., see Schober, M., 41
Fleenor, D.G., see Eldridge, J.C., 248
Freud, T.F., GABAergic septohippocampal neurons contain parvalbumin, 375

Gerhardt, G., see Eriksdotter-Nilsson, M., 269
Gerstner, G.E., Goldberg, L.J. and De Bruyne, K., Angiotensin II-induced rhythmic jaw movements in the ketamine-anesthetized guinea pig, 233
Gibbs, J., see De Saint Hilaire-Kafi, Z., 152
Ginsberg, S., see Starkstein, S.E., 16
Goldberg, L.J., see Gerstner, G.E., 233
Goldstein, M., see Kitahama, K., 156
Granholm, A.-C., see Eriksdotter-Nilsson, M., 269
Greenberg, D.A., see Marks, S.S., 196
Guevara-Aguilar, R., see Escobar, M., 368
Gunion, M.W., Taché, Y., Rosenthal, M.J., Miller, S., Butler, B. and Zib, B., Bombesin microinfusion into the rat hypothalamic paraventricular nucleus increases blood glucose, free fatty acids and corticosterone, 47

Haba, K., see Mizukawa, K., 187
Hagberg, H., see Kjellmer, I., 241
Hamada, K., see Watanabe, Y., 143

Haroutunian, V., see Anderson, J., 391
Haroutunian, V., see Benson, D.M., 399
Hayaishi, O., see Watanabe, Y., 143
Hayashi, Y., see Yajima, Y., 309
Hazlett, J.C., see Michael, D.B., 227
Heistad, D.D., see Faraci, F.M., 121
Hernandez, L., see Mifsud, J.-C., 365
Herz, A., see Ableitner, A., 326
Hetzel, F.W., see Mesina, J.E., 181
Hoebel, B.G., see Mifsud, J.-C., 365
Hoehn, K. and White, T.D., Evoked release of endogenous adenosine from rat cortical slices by K^+ and glutamate, 149
Hoffer, B., see Eriksdotter-Nilsson, M., 269
Holmes, W.R., The role of dendritic diameters in maximizing the effectiveness of synaptic inputs, 127

Indyk, E., see Britton, D.R., 205

Jami, L., Vejsada, R. and Zytnicki, D., Observations on static and dynamic responses of muscle stretch receptors in kittens, 34
Jin, J.-G., see Takaki, M., 199
Johnson Jr., E.M., Osborne, P.A. and Taniuchi, M., Destruction of sympathetic and sensory neurons in the developing rat by a monoclonal antibody against the nerve growth factor (NGF) receptor, 166
Jouvet, M., see Kitahama, K., 156
Juliano, S.L., Dusart, I. and Peschanski, M., Somatic activation of thalamic neurons transplanted into lesioned somatosensory thalamus, 356
Kaas, J.H., see Krubitzer, L.A., 161
Kaneko, T., see Watanabe, Y., 143
Kater, S.B., see Mattson, M.P., 337
Kerr, D.S., see Eldridge, J.C., 248
Kitahama, K., Okamura, H., Goldstein, M., Natgatsu, I., Bérod, A. and Jouvet, M., A new group of tyrosine hydroxylase-immunoreactive neurons in the cat thalamus, 156
Kitao, Y., see Moriizumi, T., 315
Kjellmer, I., Andiné, P., Hagberg, H. and Thiringer, K., Extracellular increase of hypoxanthine and xanthine in the cortex and basal ganglia of fetal lambs during hypoxia-ischemia, 241
Krubitzer, L.A. and Kaas, J.H., Cortical integration of parallel pathways in the visual system of primates, 161
Kudo, M., see Moriizumi, T., 315

Lagiorgia, M., see Amir, S., 361
Landau, E.M., see Benson, D.M., 399
Landfield, P.W., see Eldridge, J.C., 248
Lieberburg, I., see Anderson, J., 391
Lopes da Silva, F.H., see Boeijinga, P.H., 257
Lotstra, F., Schiffmann, S.N. and Vanderhaeghen, J.-J., Neuropeptide Y-containing neurons in the human infant hippocampus, 211

MacLean, D.B., Bennett, B., Morris, M. and Wheeler, F.B., Differential regulation of calcitonin gene-related peptide and substance P in cultured neonatal rat vagal sensory neurons, 349
Mallart, A., see Bourret, C., 403
Marks, S.S., Watson, D.L., Carpenter, C.L., Messing, R.O. and Greenberg, D.A., Calcium channel antagonist receptors in cerebral cortex from alcoholic patients, 196
Maru, E., Ashida, H. and Tatsuno, J., Long-lasting reduction of dentate paired-pulse depression following LTP-inducing tetanic stimulations of perforant path, 112
Mattson, M.P. and Kater, S.B., Excitatory and inhibitory neurotransmitters in the generation and degeneration of hippocampal neuroarchitecture, 337
Mayhan, W.G., see Faraci, F.M., 121
Mersey, J.H., see Starkstein, S.E., 16
Mesina, J.E., Page, R.H., Hetzel, F.W. and Chopp, M., Administration of L-2-oxothiazolidine-4-carboxylate increases glutathione levels in rat brain, 181
Messing, R.O., see Marks, S.S., 196
Michael, D.B., Hazlett, J.C. and Mitchell, J.A., Innervation of the pituitary gland by supraependymal neurons, 227
Mifsud, J.-C., Hernandez, L. and Hoebel, B.G., Nicotine infused into the nucleus accumbens increases synaptic dopamine as measured by in vivo microdialysis, 365
Miller, S., see Gunion, M.W., 47
Milner, T.A., Abate, C., Reis, D.J. and Pickel, V.M., Ultrastructural localization on phenylethanolamine *N*-methyltransferase-like immunoreactivity in the rat locus coeruleus, 1
Mitchell, J.A., see Michael, D.B., 227
Mizukawa, K., Takayama, H., Sato, H., Ota, Z., Haba, K. and Ogawa, N., Alterations of muscarinic cholinergic receptors in the hippocampal formation of stressed rat: in vitro quantitative autoradiographic analysis, 187
Moran, T.H., see Starkstein, S.E., 16
Moriizumi, T., Nakamura, Y., Tokuno, H., Kudo, M. and Kitao, Y., Synaptic organization of the pedunculopontine tegmental nucleus of the cat, 315
Morris, M., see MacLean, D.B., 349
Mumby, S.M., see Vogel, S.S., 281
Murakami, N. and Watanabe, T., Activaton of ACTH release is mediated by the same molecule as the final mediator, PGE_2, of febrile response in rats, 171

Nagatsu, I., see Kitahama, K., 156
Nakamura, Y., see Moriizumi, T., 315
Nakayama, S., see Takaki, M., 199
Nicolaidis, S., see De Saint Hilaire-Kafi, Z., 152
Nicoletti, F., see Clementi, G., 138

Ogawa, N., see Mizukawa, K., 187
Okamura, H., see Kitahama, K., 156
Olson, L., see Eriksdotter-Nilsson, M., 269
Ondo, J.G., see Boggan, W.O., 382
Osborne, P.A., see Johnson Jr., E.M., 166
Ota, Z., see Mizukawa, K., 187

Page, R.H., see Mesina, J.E., 181
Patrick, R.L., see Colby, K.A., 103
Peschanski, M., see Juliano, S.L., 356
Pickel, V.M., see Milner, T.A., 1
Pollock, R., see Amir, S., 361

Reis, D.J., see Milner, T.A., 1
Richter-Levin, G. and Segal, M., Raphe cells grafted into the hippocampus can ameliorate spatial memory deficits in rats with combined serotonergic/cholinergic deficiencies, 184
Roberts, J.L., see Anderson, J., 391
Robinson, R.G., see Starkstein, S.E., 16
Rogers, L.J., see Adret, P., 59
Rosenthal, M.J., see Gunion, M.W., 47

Sako, K., see Tanaka, S., 385

Sato, H., see Mizukawa, K., 187
Scapagnini, U., see Clementi, G., 138
Schiffmann, S.N., see Lotstra, F., 211
Schmechel, D.E., see Walsh, T.J., 24
Schober, M., Fischer-Colbrie, R. and Winkler, H., Ontogenesis of chromogranin A and B and catecholamines in rat adrenal medulla, 41
Schonberg, M., see Vogel, S.S., 281
Schwartz, J.H., see Vogel, S.S., 281
Segal, M., see Richter-Levin, G., 184
Seiger, Å., see Eriksdotter-Nilsson, M., 269
Seil, F.J., see Bourdette, D.N., 175
Shewey, L.M., Brot, M., Szot, P. and Dorsa, D.M., Enhanced phosphoinositol hydrolysis in response to vasopressin in the septum of the homozygous Brattleboro rat, 95
Shnayder, L., see Starkstein, S.E., 16
Snyder, S., see Anderson, J., 391
Starkstein, S.E., Ginsberg, S., Shnayder, L., Bowersox, J., Mersey, J.H., Robinson, R.G. and Moran, T.H., Developmental and hormonal factors in the sexually dimorphic, asymmetrical response to focal cortical lesions, 16
Szot, P., see Shewey, L.M., 95

Taché, Y., see Gunion, M.W., 47
Takaki, M., Jin, J.-G. and Nakayama, S., Possible involvement of calcitonin gene-related peptide (CGRP) in non-cholinergic non-adrenergic relaxation induced by mesenteric nerve stimulation in guinea pig ileum, 199
Takayama, H., see Mizukawa, K., 187
Tanaka, S., Sako, K., Tanaka, T. and Yonemasu, Y., Regional calcium accumulation and kainic acid (KA)-induced limbic seizure status in rats, 385
Tanaka, T., see Tanaka, S., 385
Taniuchi, M., see Johnson Jr., E.M., 166
Tatsuno, J., see Maru, E., 112
Thiringer, K., see Kjellmer, I., 241
Thompson, T.L., see Colby, K.A., 103
Tokuno, H., see Moriizumi, T., 315
Vanderhaeghen, J.-J., see Lotstra, F., 211
Vejsada, R., see Jami, L., 34
Vogel, S.S., Chin, G.J., Mumby, S.M., Schonberg, M. and Schwartz, J.H., G proteins in *Aplysia*: biochemical characterization and regional and subcellular distribution, 281
Von Bernhardi, R. and Alvarez, J., Is the supply of axoplasmic proteins a burden for the cell body? Morphometry of sensory neurons and amino acid incorporation into their cell bodies, 301

Wallace, W., see Anderson, J., 391
Walsh, T.J., Emerich, D.F. and Schmechel, D.E., GM1 ganglioside attenuates the behavioral deficits but not the granule cell damage produced by intradentate colchicine, 24
Watanabe, T., see Murakami, N., 171
Watanabe, Y., see Watanabe, Y., 143
Watanabe, Y., Watanabe, Y., Hamada, K., Bommelaer-Bayt, M.-C., Dray, F., Kaneko, T., Yumoto, N. and Hayaishi, O., Distinct localization of prostaglandin D_2, E_2 and F_2 binding sites in monkey brain, 143
Watson, D.L., see Marks, .S., 196
Wheeler, F.B., see MacLean, D.B., 349
White, T.D., see Hoehn, K., 149
Winkler, H., see Schober, M., 41

Yajima, Y. and Hayashi, Y., Electrophysiological evidence for axonal branching of ambiguous laryngeal motoneurons, 309
Ygge, J., Neuronal loss in lumbar dorsal root ganglia after proximal compared to distal sciatic nerve resection: a quantitative study in the rat, 193
Yonemasu, Y., see Tanaka, S., 385
Yumoto, N., see Watanabe, Y., 143

Zalc, B., see Bourdette, D.N., 175
Zhou, M., see Chiang, C.-Y., 293
Zib, B., see Gunion, M.W., 47
Zytnicki, D., see Jami, L., 34